


mgbiologics.com

RATTLERANTIVENIN

ANTIVENIN ANTIVENIN ANTIVENIN ANTIVENIN ANTIVENIN

ANTIVENIN CROTALIDAE POLYVALENT EQUINE ORIGIN



RATTLERANTIVENIN™

Shown to be effective against envenomation
due to North American Crotalidae.



PROUDLY MADE IN THE USA.

ANTIVENIN ANTIVENIN ANTIVENIN ANTIVENIN ANTIVENIN

ANTIVENIN CROTALIDAE POLYVALENT EQUINE ORIGIN

RATTLERANTIVENIN

Use: Rattler Antivenin is shown to be effective against envenomation in canine and equine due to North American Crotalidae. Crotalidae refers to cottonmouths/water moccasins, copperheads and rattlesnakes.

Use of Rattler Antivenin within 24 hours of snakebite neutralizes venom, decreases swelling, minimizes pain and decreases temperature of canine and equine patients.

Rattler Antivenin contains antivenom antibodies collected from healthy horses immunized against Western Diamondback Rattlesnake, Eastern Diamondback Rattlesnake, Prairie Rattlesnake and Mohave Rattlesnake Type A.

Administration: Administer to effect intravenously using a filtered IV set over a 20-60 minute period. Warm to body temperature using a warm water bath prior to administration. One to two 50ml doses will be sufficient in most cases regardless of body size (equine, small canine, large canine).

Administer Rattler Antivenin as is, without reconstitution or dilution. Do not mix this product with other fluids. As with any antivenin, additional fluids and supplemental therapies can be used. Rattler Antivenin has been administered in conjunction with, prior to, or immediately after other fluids and no direct risk or contraindication to the patient has been reported. Use professional judgment.

Age and Species: Canine: Safe for use eight weeks and older. Rattler Antivenin has been used safely in late-term gestation but has not been evaluated in lactating bitches. Equine: Safe for use at any age. Rattler Antivenin has not been evaluated in gestating or lactating mares.

Ingredients: Equine-derived Crotalidae Polyvalent Antivenin, Sodium Citrate

Caution: Every reasonable precaution has been taken to safeguard this product. While this serial has been reviewed and found to be safe, there is always the possibility, while rare, of adverse events. Please review your patient's history of receipt of equine protein products prior to administration.

Each box contains two 50mL doses of Rattler Antivenin.

Keep frozen below -5°C (23°F) until use.

See package insert for detailed product information.

Restricted to use by or under the direction of a licensed veterinarian.

RATTLERANTIVENIN

RATTLERANTIVENIN



Mg Biologics

Ames, IA 50014 | 877-769-2340 | mgbiologics.com

U.S.Vet Lic. No. 614